



I'm not robot



**Next**

# Benzonatate 200 mg capsule information



The symptoms of overdose can appear 15-20 minutes after swallowing the capsule, and death can happen in about one hour. This can help make sure you stay safe while you take this drug. You shouldn't break, chew, cut, or crush this drug. This drug is used to relieve coughing. This may cause you to feel dizzy or faint. If you have an allergic reaction, call your doctor or local poison control center right away. Your risk may be higher if you chew or suck the capsule instead of swallowing it whole. It starts working about 15-20 minutes after you swallow it. Don't break, chew, cut, or crush this medication. Not every pharmacy stocks this drug. A class of drugs is a group of medications that work in a similar way. Keep this drug away from light. Don't store this medication in moist or damp areas, such as bathrooms. Generic drugs usually cost less. However, because drugs affect each person differently, we cannot guarantee that this list includes all possible dosages. Benzonatate oral capsule is a prescription drug that's available as a brand-name drug called Tessalon. Keep it in your carry-on bag. Don't worry about airport X-ray machines. Keep this medication in the container it came in, tightly closed. When filling your prescription, be sure to call ahead to make sure your pharmacy carries it. Storage Store benzonatate at room temperature between 59°F and 86°F (15°C and 30°C). When flying, never put it into a checked bag. This drug starts working about 15 to 20 minutes after you swallow it. Always discuss possible side effects with a healthcare provider who knows your medical history. An interaction is when a substance changes the way a drug works. Don't take this drug again if you've ever had an allergic reaction to it. You may need to decide whether to stop breastfeeding or stop taking this medication. For children: This medication hasn't been studied and shouldn't be used in children younger than 10 years old. Some may be better suited for you than others. It may also cause negative effects on your heart and blood vessels. Psychiatric problems: This drug may cause confusion and visual hallucinations (seeing something that isn't there). Why it's used Benzonatate is used to relieve coughing. See the FDA's Safe Disposal of Medicines website ( for more information if you do not have access to a take-back program. It is important to keep all medication out of sight and reach of children as many containers (such as weekly pill minders and those for eye drops, creams, patches, and inhalers) are not child-resistant and young children can open them easily. If you are trying to access this site from the United States and believe you have received this message in error, please reach out to legal@goodrx.com and let us know. This means you may need to take it with other medications. How it works Benzonatate belongs to a class of drugs called antitussive agents. To help prevent interactions, your doctor should manage all of your medications carefully. You should always consult your doctor or other healthcare professional before taking any medication. Your dosage, drug form, and how often you take the drug will depend on your age, the condition being treated, how severe your condition is, other medical conditions you have, how you react to the first dose, forms and strengths. Generic: Benzonatate Form: oral capsule Strengths: 100 mg, 150 mg, 200 mg Brand: Tessalon Form: oral capsule Strengths: 200 mg Dosage to relieve coughing Adult dosage (ages 18 years and older) Take 100-200 mg three times per day as needed for your cough. This can be harmful or prevent the drug from working well. Serious side effects and their symptoms can include the following: allergic reactions, symptoms can include: trouble breathing, swelling or tightness in your throat, heart and blood vessel problems, such as very low blood pressure. To protect young children from poisoning, always lock safety caps and immediately place the medication in a safe location - one that is up and away and out of their sight and reach. It comes with risks if you don't take it as prescribed. If you don't take it at all: Your coughing will continue. If you take too much: You could have dangerous levels of the drug in your body. All possible dosages and drug forms may not be included here. Always speak with your healthcare provider about possible interactions with all prescription drugs, vitamins, herbs and supplements, and over-the-counter drugs that you are taking. This drug comes with several warnings. Numbness Call your doctor if you have numbness of your tongue, mouth, throat, or face that doesn't go away or gets worse. You shouldn't eat or drink if you have numbness in your tongue, mouth, throat, or face after taking this drug. Call your doctor if your cough isn't getting better. This makes you cough less after taking the drug. Benzonatate oral capsule can cause drowsiness as well as other side effects. There are other drugs available to treat your condition. Its effects last for 3 to 8 hours. Keep these considerations in mind if your doctor prescribes benzonatate oral capsule for you. General You can take benzonatate with or without food. You should swallow this drug whole. However, you should not flush this medication down the toilet. Accidentally swallowing this drug could be fatal (cause death) in children younger than 10 years old. Symptoms can appear as soon as 15 to 20 minutes after taking too much. Disclaimer: Our goal is to provide you with the most relevant and current information. Symptoms can include: trouble breathing, swelling or tightness in your throat, heart and blood vessel problems, such as very low blood pressure. Travel When traveling with your medication: Always carry your medication with you. Your doctor will write the number of refills authorized on your prescription. You should not need a new prescription for this medication to be refilled. However, because drugs affect each person differently, we cannot guarantee that this information includes all possible side effects. If they're more severe or don't go away, talk to your doctor or pharmacist. Serious side effects Call your doctor right away if you have serious side effects. Benzonatate may be used alone or with other medications. Benzonatate belongs to a class of drugs called Antitussives. It is not known if Benzonatate is safe and effective in children younger than 10 years of age. What are the possible side effects of Benzonatate? Benzonatate may cause serious side effects including: confusion, difficulty breathing, difficulty speaking, visual hallucinations, shortness of breath, tightness in the chest, and wheezing. Get medical help right away, if you have any of the symptoms listed above. The most common side effects of Benzonatate include: drowsiness, dizziness, headache, stomach upset, constipation, stuffy nose, hypersensitivity reactions, confusion, visual hallucinations, nausea, itching, skin breakouts, burning in the eyes, and chills. Tell the doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Benzonatate. The maximum dosage is 600 mg per day. Child dosage (ages 0 to 10 years) This medication hasn't been studied and shouldn't be used in children younger than 10 years old. Call your doctor right away if you become pregnant while taking this drug. Women who are breastfeeding: Benzonatate may pass into breast milk and may cause side effects in a child who is breastfed. However, because drugs interact differently in each person, we cannot guarantee that this information includes all possible interactions. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. It is very important to store this medication in a closed child-proof container and to keep it out of reach of children. If your symptoms are severe, call 911 or go to the nearest emergency room. This information is not a substitute for medical advice. This drug should be used only if the potential benefit justifies the potential risk to the fetus. Call 911 if your symptoms feel life-threatening or if you think you're having a medical emergency. If accidental ingestion occurs, seek medical attention immediately (see OVERDOSE). DESCRIPTION Benzonatate, a non-narcotic antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaoxaoctacosan-28-yl p-(butylamino) benzoate; with a molecular weight of 603.7. Benzonatate Capsules, USP contain 100 mg or 200 mg of benzonatate, USP. Benzonatate Capsules also contain: D&C Yellow No. 10, gelatin, glycerin, methylparaben sodium and propylparaben sodium. However, this article should not be used as a substitute for the knowledge and expertise of a licensed healthcare professional. Stop paying too much for your prescriptions! Thanks for visiting! GoodRx is not available outside of the United States. This medication needs to be swallowed whole. Keep this drug away from young children. This dosage information is for benzonatate oral capsule. Its effects last for 3-8 hours. Allergic reactions: This drug needs to be swallowed whole. This may cause spasms of your bronchi (the main passageway into your throat and airways) and vocal cords. The absence of warnings or other information for a given drug does not indicate that the drug or drug combination is safe, effective, or appropriate for all patients or all specific uses. It's also available as a generic drug. Store it at room temperature and away from excess heat and moisture (not in the bathroom). Unneeded medications should be disposed of in special ways to ensure that pets, children, and other people cannot consume them. These drugs are often used to treat similar conditions. Benzonatate works by numbing the receptors in your airways that cause you to cough. This may cause you to feel dizzy or faint, confusion, hallucinations (seeing or hearing something that isn't there), chest numbness, burning of your eyes. Disclaimer: Our goal is to provide you with the most relevant and current information. This serious side effect has only been seen in people who were also taking other medications. Overdose in children: Accidentally swallowing this drug could be fatal (cause death) in children younger than 10 years old. Be sure to tell your doctor about all medications, vitamins, or herbs you're taking. To find out how benzonatate oral capsule might interact with something else you're taking, talk to your doctor or pharmacist. Disclaimer: Our goal is to provide you with the most relevant and current information. Talk to your doctor if you think you're having a medical emergency. If accidental ingestion occurs, seek medical attention immediately (see OVERDOSE). The more common side effects of benzonatate oral capsule include: drowsiness, dizziness, headache, constipation, nausea, upset stomach, breakouts, nasal congestion. If these effects are mild, they may go away within a few days or a couple of weeks. The maximum dosage is 600 mg per day. Child dosage (ages 11 to 17 years) Take 100-200 mg three times per day as needed for your cough. In some cases, they may not be available in every strength or form as the brand-name version. Benzonatate also comes as an oral perle (smaller capsule). The drug information contained herein is subject to change and is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. Talk to your doctor about other drug options that may work for you. Disclaimer:

Healthline has made every effort to make information factually correct, comprehensive, and up-to-date. These issues include:Cough. Severe reactions may require intervention with vasopressor agents and supportive measures.Psychiatric EffectsIsolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking benzonatate in combination with other prescribed drugs.Accidental Ingestion And Death In ChildrenKeep benzonatate out of reach of children.Accidental ingestion of benzonatate resulting in death has been reported in children below age 10.Signs and symptoms of overdose have been reported within 15 to 20 minutes and death has been reported within one hour of ingestion. Talk to your pharmacist or contact your local garbage/recycling department to learn about take-back programs in your community. Brand name: Tessalon.Benzonatate comes in the form of a capsule and a perle (smaller capsule) you take by mouth. Always carry the original prescription-labeled box with you.Don't put this medication in your car's glove compartment or leave it in the car. They can't hurt your medication.You may need to show airport staff the pharmacy label for your medication. AllergiesBenzonatate can cause a severe allergic reaction. That means two things:Research in animals has shown adverse effects to the fetus when the mother takes the drug.There haven't been enough studies done in humans to be certain how the drug might affect the fetus.Talk to your doctor if you're pregnant or planning to become pregnant. People who chew or suck the capsule are at higher risk for allergic reactions. If your child swallows this drug, call your doctor or seek medical help right away. If your symptoms are severe, call 911 or go to the nearest emergency room right away.How to tell if the drug is working: You shouldn't cough as much. Be sure to avoid doing this when the weather is very hot or very cold. Symptoms of an overdose include:chokingbreathing problemsfeeling restlessshakinessseizuresbrain swellingstopping of your heartcoma (being unconscious for a long time)deathIf you think you've taken too much of this drug, call your doctor or local poison control center. You may report side effects to FDA at 1-800-FDA-1088.WARNINGHypersensitivitySevere hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. RefillsA prescription for this medication is refillable. Clinical monitoringYou and your doctor should monitor certain health issues. Instead, the best way to dispose of your medication is through a medicine take-back program. Your risk may be higher if you chew or suck this drug instead of swallowing it whole. Its effects last for 3-8 hours.Benzonatate may be used as part of a combination therapy. Taking it again could be fatal (cause death).Warnings for certain groupsFor pregnant women: Benzonatate is a category C pregnancy drug. Highlights for benzonatateBenzonatate oral capsule is available as both a generic drug and a brand-name drug. Always speak with your doctor or pharmacist about dosages that are right for you.Benzonatate oral capsule is typically used for short-term treatment. Once the numbness goes away, you can eat and drink. Children may be attracted to the shape and look of the liquid-filled capsules and may die if they swallow the medication.

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1

Page 1 of 1