

**Myotonic dystrophy anesthesia**

I'm not a robot   
reCAPTCHA

**Next**

## Myotonic dystrophy anaesthesia

Myotonic dystrophy after anesthesia. Myotonic dystrophy anesthesia guidelines. Myotonic dystrophy spinal anesthesia. Myotonic muscular dystrophy anesthesia considerations. Myotonic dystrophy anesthesia considerations. Myotonic dystrophy and anesthesia challenges. Myotonic dystrophy openanesthesia.org

- Mangla C, Bais K, Yarmush J. Myotonic dystrophy and anaesthetic challenges: a case report and review. Case Rep Anesthesiol. 2019;2019:4 282-305. PubMed Central Google Scholar

will 2. Morimoto Y, Mii M, Hirata T, Matayoshi H, Sakabe T. Controlled infusion of propofol for a patient with myotonic dystrophy. J Anesth. 2005;19:336-8. Article Google Scholar

3. Koyama K, Kohno M, Tsuzaki K, Kamiyama K, Morimoto Y. Bilevel Positive airway pressure therapy in a patient with myotonic dystrophy and postoperative respiratory failure: a case report. Saudi J Anaesth. 2020;14:241-3. Google Scholar

4. Masui K. Remimazolam besylate, a benzodiazepine, has been approved for general anesthesia!! J Anesth. 2020;34:479-82. Article Google Scholar

5. Wiltshire HR, Kilpatrick GJ, Tilbrook GS, Borkett KM. Phase I, ascending single dose placebo- and midazolam-controlled study to evaluate the safety, pharmacokinetics and pharmacodynamics of remimazolam (CNS 7056): Part II. Population pharmacokinetic and pharmacodynamic models and simulations. Anesth Analg. 2012;115:284-96. CAS Google Scholar

6. Doi M, Morita K, Takeda J, Sakamoto A, Yamakage M, Suzuki T. Efficacy and safety of remimazolam versus propofol for general anesthesia: a multicenter, blinded, randomized, parallel Phase IIb/III study. J Anesth. 2020;34:543-53. Article Google Scholar

7. Ahmed S, Naguib A, Tumin D, Tobias JD. Use of sugammadex in patients with myotonic dystrophy. Cardiol Res. 2018; One R, Matsuda S, Murao K, Nakamura K, Shirakawa M, Shingu K. Sugammadex use in a patient with myotonic dystrophy undergoing laparoscopic cholecystectomy. Masui. 2017;66:550-3. PubGoogle Scholar

8. Masamune T, Okawa I, Iwashita H, Oguchi T, Matsukawa T. Total intravenous anesthesia with propofol and remifentanil for a patient with myotonic dystrophy. Masui. 2009;58:620-2. PubMedia Google Scholar

will Page 2 Record of anesthesia. TOF train of four, post tetanus PTC count, bispectral BIS index, blood pressure BP, heart rate HR, systolic blood pressure SYS, diastolic blood pressure DIA, start of anaesthesia or end of anaesthesia,  $\Delta t$  between intubation or extubation, and start of surgery or end of surgery 1. J. Myotonic dystrophy type 2. Eur J Neurol 2002; 9: 441-7. Article CASE PubMed

Google Scholar

2. Ricker K, Jacobsen JF, et al. Myotonic dystrophy type 2: molecular, diagnostic and clinical spectrum. Neurology 2003; 60: 657-64. Article CAS PubMed

Google Scholar

3. Buzello W, Krieg N, Schlickewei A. Dangers of neostigmine in patients with neuromuscular disorders. Report of two cases. Br J Anaesth 1982; 54: 529-34. Article CASE PubMed

Google Scholar

4. Ranum LP, Rasmussen PF, Benzow KA, Koob MD, Day JW. Genetic mapping of a second site of dystrophy Nat Genet 1998; 19: 196-8. Article CAS PubMed

Google Scholar

5. Ricker K. Myotonic dystrophy Proximal myotonic myopathy. j neurol 1999; 246: 334-8. Article cas pubmed google scholar

6. Udd b, krahe r, Wallgren-Pettersson c, falck b, kalimo h. proximal myotonic dystrophy» a family with dominant autosomal muscular dystrophy, cataract, hearing loss and hypogonadism: myterogeneity disorder of neuromuscle 1997; 7: 217-28. Article cas pubmed google scholar

7. Thiel re. Mythonic response to suxametonio. br j anaesth 1967; 39: 815-21. Article cas pubmed google scholar

8. Orndahl g. mytonic human musculature: stimulation with depopularizing agents. It's me. mechanical recording of the effects of acetylcholine and choline. acta med scand 1962; 172: 739-51. CAS pubmed google scholar

9. Orndahl g. mytonic human musculature: stimulation with depopularizing agents. ssecond. a clinical-pharmaceutical study. acta med scand 1962; 172: 753-65. CAS pubmed google scholar

10. Orndahl g, stenberg k. Mythonic human muscles: stimulation with depopularizing agents. mechanical recording of the effects of succinildicolina, succinilmonocolina and decametonio. acta med scand suppl 1962; 389: 3-29. CAS pubmed

google scholar

11. Dundee jw. thiopentone in myotonia distrophy. curr res anesth analg 1952; 31: 257-62. CAS pubmed google scholar

12. Speedy h. exaggerated physiological responses to propofol in myotonic distrophy. br j anaesth 1990; 64: 110-2. Article cas pubmed google scholar

13. White da, smyth dg. exaggerated physiological responses to propofol in myotonic distrophy. br j anaesth 1990; 64: 758-9. Article cas pubmed google scholar

14. Aldridge lm. anesthetic problems in myotonic distrophy. a case report and a review of the aberdeen experience including 48 general anesthetics in 16 other patients. br j anaesth 1985; 57: 1119-30. Article cas pubmed google

scholar

15. Liquori cl, ricker k, moseley ml, et al. type 2 myotonic distrophy caused by a cctg expansion in the intron 1 of znf9. science 2001; 293: 864-7. Article cas pubmed google scholar

16. Savkur rs, philips av, cooper ta, et al. insulin receptor splicing in type 2. am j hum genet 2004; 74: 1309-13. Article cas pubmed google scholar

17. Wahbi k, meune c, bassez g, et al. non-compacting left ventricular in a patient with type 2. neuromuscul disord 2008; 18: 331-3. Article pubmed google scholar

18. Groh wj, groh mr, saha c, et al. electrocardiographic abnormalities and sudden death in the type 1. n engl j med 2008; 358: 2688-97. Article cas pubmed google scholar

19. Schoer og, ricker k, Schneider-Gold c, et al. sudden cardiac death in type 2 myotonic distrophy. neurology 2004; 63: 2402-4. CAS pubmed google scholar

20. Schneider-Gold c, beer m, kostler h, et al. involvement of the heart and skeletal muscle in myotonic distrophy2 (dm2): a quantitative study 31P-MRS and mri. muscle nerve 2004; 30:30:CASE

PubMed

Google Scholar

21. Flachenecker P, Schneider C, Cursiefen S, Ricker K, Toyka KV, Reiners K. Evaluation of cardiovascular autonomic function in myotonic dystrophy type 2 (DM2/PROMM). Neuromuscular Disord 2003; 13: 289-93. Pub Article Media Google School

22. Tielemans AA, van Vliet J, Jansen JB, van der Kooi AJ, Borm GF, van Engelen BGM. Gastrointestinal involvement is common in type 2 myotonic dystrophy. Neuromuscular Disorder 2008; 18: 646-9. Pub Article Med

Google Scholar

23. Auvinen S, Suominen T, Hannonen P, Bachinski LL, Krahe R, Udd B. Myotonic dystrophy type 2 found in two of the sixty-three people diagnosed with fibromyalgia. Artrite Rheum 2008; 58: 3627-31. Pub Article Media Google Scholar

24. Rudnik-Schoneborn S, Schneider-Gold C, Raabe U, Kress W, Zerres K, Schoser BG. Outcome and effect of pregnancy in type 2 myotonic dystrophy. Neurology 2006; 66: 579-80. Article CAS PubMed

Google Scholar

25. Modoni A, Silvestri G, Pomponi MG, Mangiola F, Tonali PA, Marra C. Characterization of the cognitive deficit model in myotonic dystrophy type 1. Arch Neurol 2004; 61: 1943-7. Article PubMed

Google Scholar

26. Gaul C, Schmidt T, Windisch G, et al. Subtle cognitive dysfunction in adult onset of type 1 (DM1) and type 2 (DM2) myotonic dystrophy. Neurology 2006; 67: 350-2. Article CAS PubMed

Google Scholar

27. Sansone V, Gandossini S, Cotelli M, Calabria M, Zanetti O, Meola G, Udd B, Ricker K. Report of the 84th ENMC workshop: PROMM (proximal myotonic myopathy) and other myotonic syndromes similar to myotrophy: 2nd workshop. 13-15 October 2000, Loosdrecht, Netherlands. Neuromuscular Disord 2002; 12: 306-17. 29. Mathieu J, Allard P, Potvin L, Prevost C, Begin P. A 10-year study of mortality in a cohort of patients with myotonic dystrophy. Neurology 1999; 52: 1658-62. CASE PubMedia Google Scholar

30. Meola G, Sansone V, Marinou K, et al. Proximal myotonic myopathy: a syndrome with a favorable prognosis? J Neurol Sci 2002; 193: 89-96. Article PubMed

Google Scholar

31. dal Mulino F, Class C, incrociatore H, Mall G, Giese A, Reimers CD. Cardiac involvement in proximal myotonic myopathy. Heart 1998; 79: 619-21. Google Scholar

32. Tielemans AA, Knuijt S, van Vliet J, de Swart BJ, Ensink R, van Engelen BG. Dysphagia is present but mild in type 2 myotonic dystrophy. Neuromuscular Disord 2009; 19: 196-8. Article CAS PubMed

Google Scholar

33. Milone M, Batish SD, Daube JR. Myotonic dystrophy type 2 with focal asymmetric muscle weakness and absence of electrical myotony. Muscle Nerve 2009; 39: 1383-5. Pub Article Media Google Scholar

34. Young N, Milone M, Daube JR. Electromyographic myotonia absent or unrecognized in myotonic dystrophy type 2. 2009; 72 Suppl 3: A823. Page 2 from: Anesthesia and myotonic dystrophy type 2: a series of cases Characteristics is Age at Surgery (yr) \* 166 (41, 60) Gender (n) Female 14 Female 5 Duration of symptoms (yr) \* 3 (1, 11) Procedures (n) Otorhinolaryngeal and oral 11 Or orthopaedic/pl General/urological 8 Ophthalmologic 1 Radiological 10 Symptoms/Detection (s) Muscle weakness 17 Clinical myotonia 11 Flaccid dysarthria 2 Cataract 9 Associated conditions of comorbidity (s) Diabetes mellitus 4 Cardiac conduction abnormalities 5 Cardiomyopathy 1 Preliminary aspiration pneumonia 1 \* Median (25%, 75% quartiles); n number of patients Purchase Individual access to the products is possible using the Add to Cart option on the article page. Access for 1 day (from the computer you are using) Pay-per-view content is for the exclusive use of the payee and may not be further distributed in paper or electronic form. The payee may view, download and/or print the article for personal, academic, research or educational use. Distribution of pay-per-view content is not permitted copies (electronic or otherwise) of the article.